



## Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact [support@jstor.org](mailto:support@jstor.org).

**A Partial Albino Hermit Thrush** (*Turdus aonalaschkæ pallasii*).—I have a male of this species taken in this vicinity October 27, 1885, which has the top of the head and the back light gray. Below white, the spots on the breast being fairly distinct. Tail and the primaries and secondaries fawn color.—JNO. H. SAGE, *Portland, Conn.*

**On two Abnormally Colored Specimens of the Bluebird** (*Sialia sialis*).—The United States National Museum has, through the generosity of their respective collectors, come into possession of two adult males of the common Bluebird which differ so much from the normal plumage of that species as to be worthy of special description.

No. 91,303, ♂ ad., collected at Escanaba, Mich., June 6, 1883, by Mr. Wm. Palmer, Washington, D. C., has the color of the upper parts agreeing exactly with that of more richly colored examples in the normal plumage (precisely as in No. 63,366, from Massachusetts, for example\*), the tint approaching very nearly to a pure ultramarine. The coloration of the anterior underparts, however, is quite abnormal, all of the cinnamon-colored feathers of the breast, etc., being blue beneath the surface, the feathers of the sides of the breast being chiefly or entirely blue, the latter color largely prevailing, the uniform cinnamon being confined to the middle of the breast and lower central portion of the throat. The posterior lower parts are white, however, as in true *sialis*, and not bluish, as in *mexicana*, and the bill is stout as in *sialis*, the measurements being quite normal.†

No. 107,218, collected in Baltimore Co., Maryland, March 31, 1885, by Mr. A. H. Jennings, of Baltimore, is abnormal only in the color of the upper parts, which are of a very rich uniform azure blue, almost precisely the same shade as in *S. arctica*, but even rather more greenish than in many examples of the latter species. Viewed in a particular light, many of the feathers, especially the rectrices and larger scapulars, show very regular and rather distinct darker bars, or 'water marks.' The shade of blue is quite identical with that of the *S. azurea*, the color being rather richer; but it is a notable fact that the cinnamon-color of the breast, etc., is as intense as in any specimen of normal *S. sialis*, and not of that pale ochrey tint always characteristic of *azurea*. The measurements of this specimen are as follows: Wing, 3.95; tail, 2.70; culmen, .65; tarsus, .80.

In order to show how little variation there is in this species according to locality, I present herewith measurements of several Florida specimens (three of them breeding birds) and examples from extreme northern localities. As to color, it may be remarked that none of the Florida exam-

---

\* Many examples of *S. mexicana* are not appreciably different in shade of blue from the richer colored specimens of *S. sialis*.

† [There is, and has been for many years, a specimen of *Sialia sialis* (No. 9105, ♂, Newtonville, Mass., March 10, 1868,) in the Museum of Comparative Zoölogy at Cambridge, which agrees very closely with the specimen here described by Mr. Ridgway, the whole throat being blue, and all the cinnamon-colored feathers of the breast blue beneath the surface, the blue showing prominently on the slightest disarrangement of the feathers.—J. A. ALLEN.]

ples in the National Museum collection exceed in richness of coloration the average of more northern specimens, No. 63,366, from Massachusetts, being, in fact, more intense in coloration than any of the Florida series.

## FLORIDA SPECIMENS.

Museum Number.	Sex and Age.	Locality.	Date.	Wing.	Tail.	Culmen.	Tarsus.
4920	♂ ad.	Florida (Nassau Co.)	(Summer)	4.10	2.00	.67	.80
58345	♂ "	" (Jacksonville).	Dec. 31.	4.00	2.80	.65	.78
81940	♂ "	" (Milton).	March 22.	3.95	2.70	.65	.75
89854	♂ "	" (Gainesville).	(Summer)	4.00	2.70	.65	.75
1,640, H. W. H.	♂ "	" (Cedar Keys).	Dec. 20.	3.90	2.75	.67	.78
			Average.	3.99	2.77	.66	.77

## NORTHERN SPECIMENS.

Museum Number.	Sex and Age.	Locality.	Date.	Wing.	Tail.	Culmen.	Tarsus.
61140	♂ ad.	District Columbia.	March 31.	4.05	2.75	.62	.80
82579	♂ "	" "	" 8.	4.00	2.60	.62	—
82580	♂ "	" "	April 11.	4.15	2.75	.65	.80
63366	♂ "	Massachusetts (Needham).	" 19.	3.90	2.60	.65	.80
77179	♂ "	Illinois (Cook Co.).	March 30.	3.95	2.70	.67	.80
89273	♂ "	Hudson's Bay Terr. (Moose Factory).	(Summer)	3.90	2.60	.67	.75
91303	♂ "	Michigan (Escanaba).	June 6.	3.95	2.80	.65	.75
			Average.	3.99	2.69	.65	.78

ROBERT RIDGWAY, *Washington, D. C.*